

Probability Spaces

Saravanan Vijayakumaran
sarva@ee.iitb.ac.in

Department of Electrical Engineering
Indian Institute of Technology Bombay

January 10, 2014

Probability Theory

- Helps us make decisions when faced with incomplete information
- A tool for modelling complex situations
- Applications
 - Statistical Inference
 - Communications
 - Signal Processing

What is Probability?

- Informally, a method of quantifying the degree of certainty of a situation
- Classical definition: Ratio of favorable outcomes and the total number of outcomes provided all outcomes are equally likely.

$$P(A) = \frac{N_A}{N}$$

- Relative frequency definition:

$$P(A) = \lim_{N \rightarrow \infty} \frac{N_A}{N}$$

- Axiomatic definition: A countably additive function defined on the set of events with range in the interval $[0, 1]$.
- The axiomatic definition will be used in this course

Sample Space

The first step in modelling a complex situation is to list all the possible outcomes

Definition

The set of all possible outcomes of an experiment is called the sample space and is denoted by Ω .

Examples

- Coin toss: $\Omega = \{\text{Heads, Tails}\}$
- Roll of a die: $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Tossing of two coins: $\Omega = \{(H, H), (T, H), (H, T), (T, T)\}$
- Coin is tossed until heads appear. What is Ω ?
- Life expectancy of a random person. $\Omega = [0, 120]$ years

Events

- An event is a subset of the sample space

Examples

- Coin toss: $\Omega = \{\text{Heads}, \text{Tails}\}$.
 $E = \{\text{Heads}\}$ is the event that a head appears on the flip of a coin.
- Roll of a die: $\Omega = \{1, 2, 3, 4, 5, 6\}$.
 $E = \{2, 4, 6\}$ is the event that an even number appears.
- Life expectancy. $\Omega = [0, 120]$.
 $E = [50, 120]$ is the event that a random person lives beyond 50 years.

Language of Events

Typical Notation	Language of Sets	Language of Events
Ω	Whole space	Certain event
ϕ	Empty set	Impossible event
A	Subset of Ω	Event that some outcome in A occurs
A^c	Complement of A	Event that no outcome in A occurs
$A \cup B$	Union	Event that an outcome in A or B or both occurs
$A \cap B$	Intersection	Event that an outcome in both A and B occurs
$A \cap B = \phi$	Disjoint sets	Mutually exclusive events

Which subsets must be events?

- An event is a subset of a sample space. But can all subsets of a sample space be events?
 - Yes, if the sample space is finite or countable
 - No, if the sample space is uncountable (example in next lecture)
- Let \mathcal{F} be a subset of 2^Ω consisting of all events
- If $\mathcal{F} \neq 2^\Omega$, which subsets of Ω must be there in \mathcal{F} ?
 - If we are interested in an event A , then A^c is also interesting

$$A \in \mathcal{F} \implies A^c \in \mathcal{F}$$

- If events A and B are interesting, then their simultaneous occurrence is also interesting

$$A, B \in \mathcal{F} \implies A \cap B \in \mathcal{F}$$

- These two requirements give us the following (Why?)

$$A, B \in \mathcal{F} \implies A \cup B \in \mathcal{F}$$

- They also give us $\phi \in \mathcal{F}$ if \mathcal{F} is nonempty (Why?)
- Any \mathcal{F} which satisfies these conditions is called a field
- To deal with infinite sample spaces, \mathcal{F} needs to be a σ -field

σ -fields

Definition

A collection \mathcal{F} of subsets of Ω is called a σ -field if it satisfies

- (a) $\phi \in \mathcal{F}$
- (b) if $A_1, A_2, \dots \in \mathcal{F}$, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$
- (c) if $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$

Examples

- $\mathcal{F} = \{\phi, \Omega\}$ is the smallest σ -field
- If $A \subseteq \Omega$, $\mathcal{F} = \{\phi, A, A^c, \Omega\}$ is a σ -field
- 2^{Ω} is a σ -field

Probability Measure

Definition

A probability measure on (Ω, \mathcal{F}) is a function $P : \mathcal{F} \rightarrow [0, 1]$ satisfying

(a) $P(\phi) = 0, P(\Omega) = 1$

(b) if $A_1, A_2, \dots \in \mathcal{F}$ is a collection of disjoint members in \mathcal{F} , then

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

Examples

- Coin toss: $\Omega = \{H, T\}, \mathcal{F} = \{\phi, H, T, \Omega\}$

$$P(\phi) = 0, P(H) = p, P(T) = 1 - p, P(\Omega) = 1$$

- Roll of a die: $\Omega = \{1, 2, 3, 4, 5, 6\}, \mathcal{F} = 2^\Omega, P(\{i\}) = p_i$ for $i = 1, \dots, 6$

$$P(A) = \sum_{i \in A} p_i \text{ for any } A \subseteq \Omega$$

Probability Space

Definition

A probability space is a triple (Ω, \mathcal{F}, P) consisting of

- a set Ω ,
- a σ -field \mathcal{F} of subsets of Ω and
- a probability measure P on (Ω, \mathcal{F}) .

Questions?