

1. Suppose the input X and output Y to a channel are related by $Y = \rho X + N$ where N is a zero-mean Gaussian random variable with variance σ^2 and ρ is a random variable independent of the noise. Assume that X is equally likely to be $\pm A$. Our goal is to decide on the value of X given the observation Y .
 - (a) [4 points] If ρ is the constant 1, what is the optimal decision rule and the resulting decision error probability?
 - (b) [4 points] If ρ takes values ± 1 with equal probability, what is the optimal decision rule and the resulting decision error probability?
2. [4 points] Find the maximum likelihood decision rule for the following 3-ary hypothesis testing problem where $\mu = \sqrt{2}\sigma$.

$$H_1 : Y \sim N(-\mu, \sigma^2)$$

$$H_2 : Y \sim N(0, e^2\sigma^2)$$

$$H_3 : Y \sim N(\mu, \sigma^2)$$

Hint: Sketch the density functions keeping in mind that the variances are unequal.

3. Consider the following binary hypothesis testing problem where the hypotheses are equally likely.

$$H_0 : Y \sim U \left[-\sqrt{\frac{e^2\pi}{2}}, \sqrt{\frac{e^2\pi}{2}} \right]$$

$$H_1 : Y \sim \mathcal{N}(0, 1)$$

U denotes the uniform distribution, \mathcal{N} denotes the Gaussian distribution and e is the base of the natural logarithm.

- (a) [4 points] Find the decision error probability of the rule which decides H_1 is true if $|Y| > \sqrt{\frac{e^2\pi}{2}}$ and decides H_0 is true if $|Y| \leq \sqrt{\frac{e^2\pi}{2}}$. Express your answer in terms of the Q function.
- (b) [4 points] Find the decision error probability of the optimal decision rule. Express your answer in terms of the Q function.