

# Generating Random Variables

Saravanan Vijayakumaran  
sarva@ee.iitb.ac.in

Department of Electrical Engineering  
Indian Institute of Technology Bombay

March 27, 2015

# Generating Random Variables

- Applications where random variables need to be generated
  - Simulations
  - Lotteries
  - Computer Games
- General strategy for generating an arbitrary random variable
  - Generate uniform random variables in the unit interval
  - Transform the uniform random variables to obtain the desired random variables

# Generating Uniform Random Variables

- $X \sim \mathcal{U}[a, b]$  has density function

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{for } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

- The distribution function is

$$F_X(x) = \begin{cases} 0 & x < a \\ \frac{x-a}{b-a} & a \leq x \leq b \\ 1 & x > b \end{cases}$$

- $Y \sim \mathcal{U}[0, 1]$  has distribution function

$$F_Y(x) = \begin{cases} 0 & x < 0 \\ x & 0 \leq x \leq 1 \\ 1 & x > 1 \end{cases}$$

- Given  $Y$ , can we generate  $X$ ?
- $(b - a)Y + a$  has the same distribution as  $\mathcal{U}[a, b]$

# Generating $\mathcal{U}[0, 1]$

- Computers can represent reals upto a finite precision
- Generate a random integer  $X$  from 0 to some positive integer  $m$
- Generate the uniform random variable in  $[0, 1]$  as

$$U = \frac{X}{m}$$

- The linear congruential method for generating integers from 0 to  $m$

$$X_{n+1} = (aX_n + c) \bmod m, \quad n \geq 0$$

where  $m, a, c$  are integers called the modulus, multiplier and increment respectively.  $X_0$  is called the starting value.

- For  $m = 10$  and  $X_0 = a = c = 7$ , the sequence generated is

$$7, 6, 9, 0, 7, 6, 9, 0, \dots$$

- The linear congruential method is eventually periodic

# Maximal Period Linear Congruential Generators

$$X_{n+1} = (aX_n + c) \bmod m, \quad n \geq 0$$

## Theorem

*The linear congruential sequence has period  $m$  if and only if*

- *$c$  is relatively prime to  $m$*
- *$b = a - 1$  is a multiple of  $p$ , for every prime  $p$  dividing  $m$*
- *$b$  is a multiple of 4, if  $m$  is a multiple of 4.*

## Remarks

- Having maximal period is not a guarantee of randomness
- For  $a = c = 1$ , we have  $X_{n+1} = (X_n + 1) \bmod m$
- Additional tests are needed (see reference on last slide)

# Generating a Bernoulli Random Variable

- The probability mass function is given by

$$P[X = x] = \begin{cases} p & \text{if } x = 1 \\ 1 - p & \text{if } x = 0 \end{cases}$$

where  $0 \leq p \leq 1$

- Generate a uniform random variable  $U \sim \mathcal{U}[0, 1]$
- Generate the Bernoulli random variable by the following rule

$$X = \begin{cases} 1 & \text{if } U \leq p \\ 0 & \text{if } U > p \end{cases}$$

- How can we generate a binomial random variable?

# The Inverse Transform Method

- Suppose we want to generate a random variable with distribution function  $F$ . Assume  $F$  is one-to-one.
- Generate a uniform random variable  $U \sim \mathcal{U}[0, 1]$
- $X = F^{-1}(U)$  has the distribution function  $F$

$$P(X \leq x) = P(F^{-1}(U) \leq x) = P(U \leq F(x)) = F(x)$$

## Example (Generating Exponential RVs)

$X$  is an exponential RV with parameter  $\lambda > 0$  if it has distribution function

$$F(x) = 1 - e^{-\lambda x}, \quad x \geq 0$$

How can it be generated?

# Generating Discrete Random Variables

- Suppose we want to generate a discrete random variable  $X$  with distribution function  $F$ .  $F$  is usually not one-to-one.
- Let  $x_1 \leq x_2 \leq x_3 \leq \dots$  be the values taken by  $X$
- Generate a uniform random variable  $U \sim \mathcal{U}[0, 1]$
- Generate  $X$  according to the rule

$$X = \begin{cases} x_1 & \text{if } 0 \leq U \leq F(x_1) \\ x_k & \text{if } F(x_{k-1}) < U \leq F(x_k) \text{ for } k \geq 2 \end{cases}$$

## Example (Generating Binomial RVs)

The probability mass function of a Binomial RV  $X$  with parameters  $n$  and  $p$  is

$$P[X = k] = \binom{n}{k} p^k (1-p)^{n-k} \quad \text{if } 0 \leq k \leq n$$

How can it be generated?

# Box-Muller Method for Generating Gaussian RVs

1. Generate two independent uniform RVs  $U_1$  and  $U_2$  between 0 and 1
2. Let  $V_1 = 2U_1 - 1$  and  $V_2 = 2U_2 - 1$
3. Let  $S = V_1^2 + V_2^2$ .
4. If  $S \geq 1$ , go to Step 1
5. If  $S < 1$ , let

$$X_1 = V_1 \sqrt{\frac{-2 \ln S}{S}}, \quad X_2 = V_2 \sqrt{\frac{-2 \ln S}{S}}$$

6.  $X_1$  and  $X_2$  are independent standard Gaussian random variables

## Proof

- $(V_1, V_2)$  represents a random point in the unit circle
- Let  $V_1 = R \cos \Theta$  and  $V_2 = R \sin \Theta$
- $\Theta \sim \mathcal{U}[0, 2\pi]$  and  $R^2 = S \sim \mathcal{U}[0, 1]$ .  $\Theta$  and  $S$  are independent
- $X_1 = \sqrt{-2 \ln S} \cos \Theta$  and  $X_2 = \sqrt{-2 \ln S} \sin \Theta$
- $X_1, X_2$  also are in polar coordinates with radius  $R' = \sqrt{-2 \ln S}$  and angle  $\Theta$

## Proof Continued

- The probability density function of  $R'$  is  $f_{R'}(r) = re^{-r^2/2}$

$$\Pr [R' \leq r] = \Pr [\sqrt{-2 \ln S} \leq r] = \Pr [S \geq e^{-r^2/2}] = 1 - e^{-r^2/2}$$

- The joint probability distribution of  $X_1$  and  $X_2$  is given by

$$\begin{aligned} P(X_1 \leq x_1, X_2 \leq x_2) &= \int_{\{(r, \theta) | r \cos \theta \leq x_1, r \sin \theta \leq x_2\}} \frac{1}{2\pi} e^{-\frac{r^2}{2}} r \, dr \, d\theta \\ &= \frac{1}{2\pi} \int_{\{x \leq x_1, y \leq x_2\}} e^{-\frac{x^2+y^2}{2}} \, dx \, dy \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_1} e^{-\frac{x^2}{2}} \, dx \cdot \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_2} e^{-\frac{y^2}{2}} \, dy \end{aligned}$$

- This proves that  $X_1$  and  $X_2$  are independent and have standard Gaussian distribution

# Acceptance-Rejection Method

- Suppose we want to generate a random variable  $X$  having density  $f$
- Suppose  $X$  is difficult to generate using the inversion method
- Suppose there is a random variable  $Y$  with density  $g$  which is easy to generate
- For some  $c \in \mathbb{R}$ , suppose  $f$  and  $g$  satisfy

$$\frac{f(y)}{cg(y)} \leq 1 \text{ for all } y.$$

- Generate a uniform random variable  $U \sim \mathcal{U}[0, 1]$
- Generate the random variable  $Y$
- If  $U \leq \frac{f(Y)}{cg(Y)}$ , set  $X = Y$ . Otherwise, generate another pair  $(U, Y)$  and keep trying until the inequality is satisfied
- To show that the method is correct, we have to show that

$$P\left(Y \leq x \mid U \leq \frac{f(Y)}{cg(Y)}\right) = F(x)$$

where  $F(x) = \int_{-\infty}^x f(t) dt$

# Example of Acceptance-Rejection Method

- Suppose we want to generate a random variable  $X$  with probability density function

$$f(x) = 20x(1 - x)^3, \quad 0 < x < 1$$

- We need a pdf  $g(x)$  such that  $\frac{f(x)}{g(x)} \leq c$  for some  $c \in \mathbb{R}$
- Consider  $g(x) = 1$  for  $0 < x < 1$

$$\frac{f(x)}{g(x)} = 20x(1 - x)^3 \leq 20 \cdot \frac{1}{4} \cdot \left(\frac{3}{4}\right)^3 = \frac{135}{64}$$

- Let  $c = \frac{135}{64} \implies \frac{f(x)}{cg(x)} = \frac{256}{27} x(1 - x)^3$
- $X$  can now be generated as follows
  1. Generate  $U \sim \mathcal{U}[0, 1]$  and  $Y \sim \mathcal{U}[0, 1]$
  2. If  $U \leq \frac{256}{27} Y(1 - Y)^3$ , set  $X = Y$
  3. Otherwise, return to step 1

## Reference

- Chapter 3, *The Art of Computer Programming, Seminumerical Algorithms (Volume 2)*, Third Edition, Pearson Education, 1998.